

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12Q 1/34</b>	<b>A1</b>	(11) International Publication Number: <b>WO 95/34678</b> (43) International Publication Date: <b>21 December 1995 (21.12.95)</b>
<p>(21) International Application Number: <b>PCT/US95/07207</b></p> <p>(22) International Filing Date: <b>7 June 1995 (07.06.95)</b></p> <p>(30) Priority Data: <b>08/259,409</b>      <b>14 June 1994 (14.06.94)</b>      <b>US</b></p> <p>(71) Applicant: <b>AMERICAN CYANAMID COMPANY [US/US];</b> <b>One Cyanamid Plaza, Wayne, NJ 07470-8426 (US).</b></p> <p>(72) Inventor: <b>SILVERMAN, Sanford, Jay; P.O. Box 657, Roosevelt, NJ 08555 (US).</b></p> <p>(74) Agents: <b>WEBSTER, Darryl, L.; American Cyanamid Company, One Cyanamid Plaza, Wayne, NJ 07470 (US) et al.</b></p>	<p>(81) Designated States: <b>AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).</b></p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: <b>SCREEN FOR INHIBITORS OF CHITINASE</b></p> <p>(57) Abstract</p> <p>The present invention is directed to a method for screening samples for the identification of agents exhibiting potential fungicidal and insecticidal activity for a wide variety of agricultural, medical and pharmaceutical uses. The method utilizes cells that comprise a plasmid-born CTS gene of <i>Saccharomyces cerevisiae</i>, which allows for over expression of chitinase. Compounds that inhibit a hydrolytic action on methyl-umbelliferyltriacyetyl chitotriose, but which are not toxic to the cells are detected by decrease in conversion of substrate.</p> <p style="text-align: right;"><b>FILE COPY</b></p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## SCREEN FOR INHIBITORS OF CHITINASE

Field of the Invention

The present invention relates to a method of screening compounds that inhibit chitinase. More specifically, the present invention relates to the identification of antifungal, insecticidal and antiparasitic compounds for use in agricultural and pharmaceutical applications.

BACKGROUND OF THE INVENTION

The polysaccharide chitin is a structural cell wall component of most fungi and is the most abundant organic skeletal component of invertebrates, making up, for example, from about 25 to 60% of the dry weight of insect cuticles. Chitin consists primarily of linear polymers of the amino sugar N-acetyl-D-glucosamine joined in 1,4- $\beta$ -glucosidic linkage. Thus, chitin bears a close resemblance to cellulose, the major structural polysaccharide of plants, the only chemical difference being that in chitin the hydroxyl group on the 2-position is an acetoamido group instead of an hydroxyl group. However, because of its widespread occurrence in fungi and arthropods, the total world-wide production of chitin vastly exceeds cellulose.

Many fungi and arthropods having chitinous cell walls or exoskeletons are injurious to plants and animals, causing a legion number of diseases including, but not limited to, wheat eyespot, rice sheath blight, damping off, apple scab, pepper

- 2 -

botrytis, rice blast, sugar beet cercospora, tomato  
early blight, wheat leaf rust, and wheat powdery-  
mildew. Fungal species also cause a myriad of  
cutaneous and systemic mycoses in human beings and  
5 other animals, including, but not limited to,  
candidiasis, histoplasmosis, blastomycosis,  
pneumocystis, sporotrichosis and cryptococcosis.  
Insects can act as vectors of viruses causing  
arboviral encephalitides, yellow fever, and dengue,  
10 protozoa causing malarias, trypanosomiasis, and  
leishmaniases, and various harmful helminths.  
Crustaceans also carry some infectious helminths and  
trematodes.

Most fungicides and insecticides that are  
15 used to control or cure these diseases by killing or  
controlling their causative agents, intermediate  
hosts, or vectors employ various modes of action  
including physical poisons that suffocate or  
desiccate organisms; protoplasmic poisons such as  
20 arsenicals that kill by precipitating or  
deactivating proteins, enzymes or other cellular  
constituents; respiratory poisons that deactivate  
respiratory enzymes; and various poisons that affect  
different tissue systems such as tubules or nerves.  
25 Of course, preferred agents do not injure the host  
plant or animal, and most preferably have no effect  
whatsoever on the host. Because of the complexity  
and interdependence of life processes, however,,  
this goal is not always achieved, so that many  
30 fungicides and insecticides exhibit some toxicity to  
the host. Others cause unexpected side effects.

Since chitin is not a usual constituent of  
most plants and vertebrates, chitin biosynthesis  
inhibitors can be employed as selective antifungal  
35 and/or insecticide agents. Applied to ornamental or

- 3 -

edible plants or animals, these offer the advantage of targeting undesirable fungi or insects without harming significantly the host plant or vertebrate animal. While much attention has been paid to chitin synthesis, there have been very limited studies targeting screens which exploit chitin degradation. 1-(2,6-Dichlorobenzoyl)-3-(3,4-dichlorophenyl)urea, for example, has been suggested as a chitin-inhibiting insecticide. Antifungals that have also been found to inhibit chitin synthesis include nikkomycin and polyoxin D.

#### SUMMARY OF THE INVENTION

It is an object of the present invention to provide a screening test for the identification of agents exhibiting potential fungicidal and insecticidal activity for a wide variety of agricultural, medical and pharmaceutical uses. This and other objects are accomplished by the present invention, which is directed to a method of screening compounds useful as antifungal, insecticidal and antiparasitic agents. The preferred method utilizes cells that comprise a plasmid-born CTS gene of *Saccharomyces cerevisiae* which allows for over expression of chitinase. Compounds that inhibit a hydrolytic action on methyl-umbelliferyltriacyl chitotriose, but which are not toxic to the cells, are detected by decrease in conversion of substrate. The present invention allows for the high volume screening of chemicals and fermentations for inhibitors of chitinase for insecticide, antiparasitic and fungicide applications.

- 4 -

DETAILED DESCRIPTION OF THE INVENTION

5           The discovery of most pharmaceutical and agricultural chemical products has resulted, at least in part, from the screening of either chemical libraries or natural products. A variety of screening systems that employ mammalian cells or yeast cells are well-known by those skilled in the art and have been described.

10           In the practice of this invention, test samples are incubated in the presence of cultures of any fungal species that produces chitin, such as unicellular fungi. A preferred method employs common baker's yeast, *Saccharomyces cerevisiae*,  
15           because it is readily available and easy to culture.

          A preferred method comprises adding a test sample to a *Saccharomyces cerevisiae* culture. The test sample is introduced to a disk or a well on a culture plate in a standard diffusion assay using  
20           solidified media, or introduced into one of a series of equivalent tissue culture tubes or bottles in a standard turbidity assay using liquid media. The culture is incubated for such time under such conditions sufficient to observe yeast cell growth  
25           inhibition in a corresponding culture or culture plate area. The extent of growth of the culture containing or surrounding the test sample is compared with the extent of growth in the culture or culture area containing no test sample. The extent  
30           of toxicity of the test sample is determined by observing whether growth in the presences of test sample is substantially the same as growth in its absence.

35           The present invention relies on the expression of chitinase on the yeast cell surface

- 5 -

and detection of enzyme activity with a substrate by observing the extent of substrate conversion.

Any type of substrate for chitinase that affords easy detection of enzyme activity can be employed including fluorescent substrates, colored dyes and turbid substrates that become clear when exposed to enzyme activity. Examples of substrants include: colloidal chitin, glycol chitin, 3-4 dinitrophenyl tetra-N-acetyl chitotetra oside, 4-methylumbelliferyl di-N-acetyl chitobioside, 4-methylumbelliferyl tri-N-acetyl chitotrioside, or 4-methylumbelliferyl tetra-N-acetyl chitotetraoside.

Any type of solidified or liquid media that will support growth and reproduction of *S. cerevisiae* may be employed as cultures in the method of this invention. Numerous yeast media are known to the skilled artisan, and include, for example, yeast synthetic dextrose (SD) containing glucose, vitamins, minerals, and water. Preferred media are solidified by adding agar or gelatin; especially preferred are media solidified with agar.

Growth in solidified cultures is ordinarily observed visually as turbid areas of growth around disks or wells in the culture plate. Growth in liquid cultures is observed visually, but is ordinarily determined spectrophotometrically as enhanced optical density (OD) at about 550 to 650 nm.

A distinct advantage of the invention is its speed and simplicity. Baker's yeast is readily available and inexpensive. Using solidified media in culture plates, the protocol is extremely simple. Many samples can be readily analyzed in a short time.

It is another advantage of the invention

- 6 -

that only small amounts of biochemical or chemical agents are required in the test. In a standard assay, for example, which employs solidified media in a plate, as little as 20  $\mu$ g of a biochemical or chemical test sample can be applied to a disk or in a well.

In a preferred embodiment of this invention, the screen relies on the expression of excess chitinase on the cell surface and direct detection of enzyme activity with a fluorescent substrate. This embodiment is achieved by using yeast cells transformed with a plasmid carrying the chitinase gene. The transformed yeast cells are dispersed and grown in agar, substrate is applied and the extent of the reaction is assayed by observation of the plate under ultraviolet illumination.

In a preferred embodiment of the invention, the fluorescent substrate is 4-methylumbelliferyl-tri-N-acetyl chitotrioxide (MUC).

In a preferred embodiment of the invention, the chitinase gene is carried on plasmid pCT21 which is essentially the same as plasmid pCT3 (Kuranda, M.J. and Robbins, P.W., (1987) Proc. Natl. Acad. Sci. USA vol. 84, 2585-2589) except that the vector sequences comprise the well known and commercially available YEp24 instead of the described YEAp24. pCT21 over-expressed chitinase from 4-25 fold over wild type cells lacking the plasmid.

In a particularly preferred embodiment, the present invention comprises a primary screen and secondary screen which increases both sensitivity and specificity of the invention. It should be understood by those skilled in the art that the



- 7 -

present invention can be practiced using the primary screening alone, the secondary screen alone or a combination of the primary screen and secondary screen. One skilled in the art should also appreciate that the primary screen could be practiced after the secondary screen depending on the ultimate objective of the invention.

Standard in vitro and in vivo fungicide and insecticide discovery screens are employed as tertiary tests to prioritized actives from the present primary screen and the secondary screen. These in vitro screens test samples for their ability to inhibit the growth of selected phytopathogenic fungi cultured in nutrient agar (or insects). These include fungi causing wheat eyespot (*Pseudocercospora herpotrichoides*), rice sheath blight (*Rhizoctonia solani*) and damping off (*Fusarium oxysporum*); all synthesize chitin-containing cell walls.

In in vivo screens, a variety of phytopathogenic fungi are used to infect plants treated with test compounds. Active compounds block or reduce the appearance of disease symptoms. A number of model plant infections are employed in the screen and include chitin-containing fungi that cause apple scab (*Venturia inaequalis*), pepper botrytis (*Botrytis cinerea*), rice blast (*Pyricularia oryzae*), sugar beet cercospora (*Cercospora beticola*), tomato early blight (*Alternaria solani*), wheat leaf rust (*Puccinia recondita tritici*), and wheat powdery mildew (*Erysiphe graminis tritici*). The most potent test compounds in these assays are active in the 10 ppm range.

The following examples are presented to further illustrate and explain the present invention

- 8 -

and should not be taken as limiting in any regard.

#### Examples

5      **Example 1 - PREPARATION OF YEAST EXTRACTS (Chitinase extract)**

Inoculate 200 ml and grow EC18-4B pCT as for primary screen. Centrifuge culture at 300rpm for 10 min. Pour off supernatant, re-spin 1 min and remove remaining supernatant. Resuspend cells in 4  
10 X 1ml of buffer (25mM Mes-0.1% digitonin-0.1%2-mercaptoethanol, pH 6.3, for chitinase; 0.1M Na citrate, pH 5.0 for glucanase) in a screw cap centrifuge tube. Add approximately 0.5mm glass  
15 beads to just below the meniscus and Vortex vigorously or put on a Mini-Bead-Beater for approximately 2 minutes. Remove liquid, wash beads with 0.5ml buffer, add wash to the rest of the extract. Centrifuge at 10,000 rpm for 2 minutes.  
20 Remove supernatant, measure its volume and add 1/2 volume glycerol and vortex gently. Store in aliquots at -80°C (the extracts are then tested in the appropriate enzyme assay to ensure sufficient activity and linearity over time. 5 µl and 45  
25 minutes is standard) at 30°C.

**Example 2 - PRIMARY IN VITRO ENZYME ASSAY**

Grow the cell to stationary phase (overnight) in SD plus supplements without agar and without chloramphenicol. The Uracil requirement is  
30 supplied by the chitinase gene-containing plasmid pCT21. To retain the plasmid uracil must NOT be added. Cells can be stored for one week or more after suspending them in the agar medium and pouring  
35 the plate. For large scale, high capacity screening

- 9 -

add sterile glycerol (15% final volume) to the fresh, stationary phase cultures and store at  $-80^{\circ}\text{C}$  in aliquots appropriate for test plate volume.

5 If using frozen cells, allow to thaw at room temperature and vortex. Add cells at 1:50 to SD plus supplements (with chloramphenicol when testing and natural products) that has been melted and cooled to about  $50^{\circ}\text{C}$ . Allow to solidify and apply compounds with a Clonemaster, or cut wells and add fermentations robotically. The positive control is allosamidin ( $1.25\mu\text{g}$ ).

Allow cells to grow 24 hours at  $30^{\circ}\text{C}$ .

15 Melt the assay overlay agar, cool to about  $50^{\circ}\text{C}$ , and add MUC. Mix well and apply to plate situated on a level surface for even distribution of soft agar. A large square plate that has 150 ml of agar requires 330 ml of overlay.

When plates have cooled, place them back at  $30^{\circ}\text{C}$  and score after about two hours.

20 Alternatively, MUC can be added directly to the SD plus supplements at the same time cells are added. In this case, no assay overlay agar is necessary.

25 While wearing UV-protective glasses, place plates on a UV transilluminator (cover off, agar face down) and observe level of blue fluorescence around wells. Mark wells with decreased fluorescence (compare to allosamidin). In normal light, determine whether decrease in fluorescence is due to growth inhibition. Score a compound positive that decreases fluorescence but does not inhibit growth.

30 To mimic chitinase inhibition, various high-pH solutions (which are known to inhibit chitinase) are applied to filter discs and placed on

35

- 10 -

the agar. Good fluorescence is obtained after overlaying substrate, and a zone of inhibition is seen only with the strongest alkali solutions tested (e.g. 10M NaOH). Other material designed to mimic fermentations and the potential interfering compounds therein (proteases [e.g. 10 mg/ml Proteinase K, trypsin], buffering capacity) are found not to cause a false positive signal. Antimicrobial test agents are unnecessary in this screen since antifungals and other compounds that nonspecifically inhibit growth are scored as negative. Allosamidin, an analog of the natural substrate chitin, is known to inhibit *Bombyx mori* and *Saccharomyces* chitinase, but not plant chitinases (e.g. the enzyme from yam). Allosamidin is very clearly positive at about 1.25-2.5 $\mu$ g per application. However, the more recently found compound, demethylallosamidin has been shown to be 100 fold more active than allosamidin and should be observable at 0.0125 $\mu$ g per application (500ng/ml). Applying the compound at a high concentration in DMSO on the agar surface as is done for other high-volume plate screens works as well as filter disk application. The best results for "welled" plates is to apply the compounds in 25 $\mu$ l in the well. In a reconstruction experiment 10 randomly chosen fermentation broths are "spiked" with 5  $\mu$ g allosamidin in 25 $\mu$ l and applied to a primary screen plate. All 10 wells are clearly positive, whereas the same broths without allosamidin are negative.

The secondary assay is much more sensitive. Allosamidin inhibits chitinase to approximately the 50% level at 100ng/ml final concentration in the assay. No inhibition of glucanase is detectable.

- 11 -

The screen is tested with thirty four standard test compounds exhibiting a variety of modes of action as well as with seventy six compounds that comprise a standard panel of antibiotics, as disclosed in Tables 1 and 2. All were negative, as are fermentation 44D048, aristeromycin, diflubenzuron (dimilin) and known chitin synthase inhibitors polyoxin and nikkomycin.

For chemical bank screening approximately 0.03% of compounds are judged positive in the primary screen. After processing over 20,000 chemicals, none have passed the secondary screen. For natural products the primary screen results in 0.9% active, but in over 15,000 assays only 2 (0.013%) have passed the secondary screen.

*Table 1 - Standard Fungicide Panel*

<u>Compound</u>	<u>Target</u>
Amphotericin B	plasma membrane (polyene)
Cerulenin	fatty acid biosynthesis
Haloprogin	respiration
ketoconazole	ergosterol biosynthesis (lanosterol 14a-demethylase)
iconazole	ergosterol biosynthesis (lanosterol 14a-demethylase)
diniconazole	ergosterol biosynthesis (lanosterol 14a-demethylase)
econazole	ergosterol biosynthesis (lanosterol 14a-demethylase)
fenarimole	ergosterol biosynthesis (sterol d14 reductase)
tridemorph	ergosterol biosynthesis (sterol d14 reductase)
tolnaftate	ergosterol biosynthesis (squalene monooxygenase)
U18666A-	ergosterol biosynthesis (squalene cyclase)
cycloheximide	protein biosynthesis
polyoxin D	chitin biosynthesis (cell wall)
nikkomycin	chitin biosynthesis (cell wall)
nocodazole	microtubule
benomyl	microtubule
maneb	multi-target
metalaxyl	rRNA biosynthesis
vinclozolin	lipid peroxidation
kanamycin	mitochondria
tunicamycin	glycoprotein biosynthesis
carboxin	succinate dehydrogenase
cyanobuturate	microtubule (plant)
antimycin	respiration
glyphosate	herbicide (aromatic amino acid biosynthesis)
phosphinothricin	herbicide (glutamine biosynthesis)
aminotriazole	herbicide (histidine biosynthesis)
sulfometuron methyl	herbicide (branched chain amino acid biosynthesis)
pendimethalin	herbicide (microtubule)

*Table 2 - Standard Antibiotic Panel*

pimaricin (tonnecetin)	streptogramin ("type")
monazomycin	nystatin
aspartocin	bacitracin
clavacin	citinin
avoparcin	isoquinocycline
neutamycin	A1531
leucomycin	AO341a
angustmycin A & C	gliotoxin
gibberellic acid	puromycin
puromycin aminonucleoside	BM123a
etamycin	mocimycin
neomycin	viomycin
netropsin	lincomycin
picromycin	A9537
AN272a	levomycin
AM374	antiprozoin
BL580 zeta	actithiazic acid
hamycin	carbomycin
frenolicin	fusarinic acid
BL580a	tylosin
declomycin	tetrahydro spiramycin
usnic acid	geldanamycin
Z122OA	BM782a
BO2964 complex	chloramphenicol
A8363	actinomycin
BM123a	AD97
phenazine a	paromomycin
streptomycin	A4825
BO2964 complex	nucleocidin
nonactin	valinomycin
C19004 complex	avilamycin
V214W	V214X
vancomycin	ristocetin
relomycin	CO8078a
blastocidinS	4-dedimethylamino-4-methylamino-anhydro-tetracycline

- 14 -

Example 3 - Secondary in vitro Enzyme Assay

Actives from the primary screen are put through the secondary assay which takes about 2 hours to perform for several dozen samples, including preparation and analysis. This consists of two different in vitro enzyme assays: glucanase and chitinase (the target). Broths that inhibit chitinase but not glucanase in these tests are judged positive. Those that display an absolute differential for these enzymes are first priority leads. Broths that show greater than twice the inhibition of chitinase versus glucanase should be followed up with secondary priority.

For each primary screen positive two enzyme assays are performed in microtiter dish wells. For each set of assays, make a Mix of buffer, Substrate and water, and add fermentation broth and enzyme:

Chitinase: Mix=1M Na citrate: 1/4 MUC:  
water, 5:10:20

Per assay:

Mix 35 $\mu$ l

broth or water (control) 10 $\mu$ l

Chitinase extract 5 $\mu$ l

Incubate for 45 min at 30°C, then add 100 $\mu$ l glycine-NaOH, mix.

Glucanase: Mix = 1M Na citrate:MUG:water,  
5:10:20

Per assay:

Mix 35 $\mu$ l

broth or water (control) 10 $\mu$ l

Chitinase extract 5 $\mu$ l

Incubate as for chitinase. Read both assays in a fluorimeter (350nm excitation, 440nm emission). Compare controls for each to broth



- 15 -

effect on each. Broths that inhibit chitinase at least twice as much as they inhibit glucanase are judged positive.

- 16 -

What is claimed:

1. A method of screening test samples for chitin degradation including a primary in vitro enzyme assay and a secondary in vitro enzyme assay, said primary assay comprising the steps of:
- (a) adding the test sample to a chitinase producing yeast culture, said culture containing a substrate for chitinase;
  - (b) incubating said test sample in said culture under conditions sufficient to detect inhibition of chitinase enzyme activity;
  - (c) comparing the extent of substrate conversion in the area around test sample with the extent of substrate conversion in the area void of said test sample;
  - (d) determining the presence of chitinase inhibition by observing the extent of substrate conversation; and
- said secondary assay comprising the steps of:
- (e) adding a test sample that exhibits chitinase inhibition in said primary assay to a series of secondary enzyme assays, said secondary enzyme assays selected from the group consisting of a chitinase assay and at least one other control enzyme assays;
  - (f) incubating said test sample in said secondary enzyme assays under conditions sufficient to detect inhibition of chitinase and

- 17 -

inhibition of said control enzyme assay;

(g) simultaneously comprising chitinase inhibition and control enzyme inhibition; and

(h) determining whether the extent of chitinase inhibition is greater than the control enzyme inhibition.

2. The method of Claim 1 wherein the yeast culture overproduces chitinase.

3. The method of Claim 2 wherein the yeast culture comprises plasmid pCT21.

4. The method of Claim 1 wherein the enzyme activity is detected with a fluorescent substrate.

5. The method of Claim 1 wherein the enzyme activity is detected with a colored dye substrate.

6. The method of Claim 1 wherein the enzyme activity is detected with a clearing of turbidity.

7. The method of Claim 1 wherein the substrate is selected from the group of consisting of colloidal chitin, glycol chitin, 3-4 dinitrophenyl tetra-N-acetyl chitotetra oside, 4-methylumbelliferyl di-N-acetyl chitobioside, 4-methylumbelliferyl tri-N-acetyl chitotrioside, or 4-methylumbelliferyl tetra-N-acetyl chitotetraoside.

8. The method of Claim 1 wherein the substrate is 4-methylumbelliferyl tri-N-acetyl chitotrioside, or 4-methylumbelliferyl tetra-N-acetyl chitotetraoside.


- 18 -

9. The method of Claim 1 wherein a positive control is selected from the group consisting of allosamidin or demethylallosamidin.
- 5 10. The method of Claim 1 wherein the enzyme assay control of step (a) is glucanase.

## INTERNATIONAL SEARCH REPORT

i. national application No.

PCT/US 95/07207

A. CLASSIFICATION OF SUBJECT MATTER		
IPC6: C12Q 1/34 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: C12Q		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search: (name of data base and, where practicable, search terms used)		
WPI, IFIPAT, CA, MEDLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	THE JOURNAL OF ANTIBIOTICS, Volume XL, No 12, May 1987, PENELOPE J. B. SOMERS ET AL, "Method for the detection and quantitation of chitinase inhibitors in fermentation broths; Isolation and insect life cycle effect of A82516" page 1751  -----	1-10
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
29 Sept 1995		24. 10. 95
Name and mailing address of the International Searching Authority		Authorized officer
 European Patent Office, P.B. 5818 Patentlaan 2 NL-2230 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016		Carolina Gómez Lagerlöf

BEST AVAILABLE COPY